

Formulas (you need very few of these!!)

| | | | |
|---|--|--|--|
| $x = x_0 + v_0t + \frac{1}{2}a_x t^2$ | $\sum \vec{F} = m\vec{a}$ | $W = \int_{\vec{x}_i}^{\vec{x}_f} \vec{F}(x) \cdot d\vec{x}$ | $T_c = \frac{5}{9}(T_f - 40) + 40$ |
| $\vec{A} \cdot \vec{B} = A B \cos\theta$ | $\vec{a} \times \vec{b} = a b \sin\phi \perp$ to both | | $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ |
| $\sin\theta = \frac{o}{h}$ | $\cos\theta = \frac{a}{h}$ | $\tan\theta = \frac{o}{a}$ | use, $g = 10 \frac{m}{sec^2}$ |
| $\tan 45^\circ = 1$ | $\sin 30^\circ = 0.5 = \cos 60^\circ, \cos 30^\circ = 0.866 = \sin 60^\circ$ | | |
| $\sin 45^\circ = 0.707 = \cos 45^\circ$ | $C = 2\pi r$ | $A = \pi r^2$ | |
| $V = \frac{4}{3}\pi r^3$ | $A_{cylinder} = 2\pi r h$ | $A_{sphere} = 4\pi r^2$ | $T_c = \frac{5}{9}(T_f - 40) + 40$ |
| $\Delta L = L\alpha\Delta T$ | $\Delta V = V\beta\Delta T$ | $T_K = T_C + 273$ | $T_F = \frac{9}{5}T_C + 32^\circ$ |
| $W = \int_{V_i}^{V_f} P dV$ | $\Delta E_{int} = Q + W$ | $Q = cm(T_f - T_i)$ | |
| $Q = Lm$ | $H = \frac{Q}{t} = kA \frac{T_h - T_c}{L}$ | | $P_r = \sigma\epsilon AT^4$ |
| $N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$ | $pV = nRT$ | $pV^\gamma = const$ | $\bar{K} = \frac{3}{2}kT$ |
| $W_{isothermal} = nRT \ln\left(\frac{V_f}{V_i}\right)$ | $v_{rms} = \sqrt{\frac{3RT}{M}}$ | | $k = 1.38 \times 10^{-23} \text{ J / K}$ |
| $R = 8.31 \text{ J/mol K}$ | $\gamma = \frac{C_p}{C_v}$ | | $C_p = C_v + R$ |
| $C_v = \frac{3}{2}R$ (ideal monatomic gas) | $E_{int} = nC_v T$ | | $\Delta E_{int} = nC_v \Delta T$ |
| $\Delta S = S_f - S_i = \int_i^f \frac{dQ}{T}$ | $\eta = \frac{ W }{ Q_h }$ | $K = \frac{ Q_c }{ W }$ | $\eta = 1 - \frac{ Q_c }{ Q_h } = 1 - \frac{T_c}{T_h}$ |
| $K = \frac{ Q_c }{ Q_h - Q_c } = \frac{T_c}{T_h - T_c}$ | | | |

| | | | | | |
|---|--|--|---|---|--|
| $\omega_0 = \frac{1}{\sqrt{LC}}$ | $\Phi = \oint \vec{E} \cdot d\vec{A}$ | $\epsilon_0 \Phi = q_{enc}$ | $\oint \vec{E} \cdot d\vec{A} = \frac{q_{enc}}{\epsilon_0}$ | | |
| $\vec{F} = \frac{q_1 q_2}{4\pi\epsilon_0 r^2} \hat{r}$ | $\vec{E} = \frac{\vec{F}}{q_o}$ | $\vec{E} = \frac{ q }{4\pi\epsilon_0 r^2} \hat{r}$ | $d\vec{E} = \frac{ dq }{4\pi\epsilon_0 r^2} \hat{r}$ | | |
| $q = \lambda L$ | $dq = \lambda ds$ | $q = \sigma A$ | $dq = \sigma dA$ | $q = \rho V$ | $dq = \rho dV$ |
| $\epsilon_0 = 8.85 \times 10^{-12} \text{C}^2 / \text{N m}^2$ | $k_e = \frac{1}{4\pi\epsilon_0} = 8.99 \times 10^9 \text{Nm}^2 / \text{C}^2$ | $\mu_0 = 4\pi \times 10^{-7} \frac{\text{Tm}}{\text{A}}$ | | | |
| $V = -\vec{E} \cdot \vec{d}$ | $\Delta U = q\Delta V$ | $V = \frac{q}{4\pi\epsilon_0 r}$ | $E_x = -\frac{\partial V}{\partial x}$ | $E_y = -\frac{\partial V}{\partial y}$ | $E_z = -\frac{\partial V}{\partial z}$ |
| $C = \frac{q}{V}$ | $C = \frac{\epsilon_0 A}{d}$ | $C_{eq} = \sum_{j=1}^n C_j$ | $\frac{1}{C_{eq}} = \sum_{j=1}^n \frac{1}{C_j}$ | | |
| $U = \frac{q^2}{2C} = \frac{1}{2} CV^2$ | $C_{eq} = \kappa C_{air}$ | $U_B = \frac{1}{2} Li^2$ | $u_B = \frac{B^2}{2\mu_0}$ | | |
| $i = \frac{dq}{dt}$ | $i = \int \vec{J} \cdot d\vec{A}$ | $R = \frac{V}{i}$ | $\rho = \frac{1}{\sigma} = \frac{E}{J}$ | $R = \frac{\rho L}{A}$ | |
| $R = R_0[1 + \alpha(\Delta T)]$ | $P = iV$ | $P = i^2 R = \frac{V^2}{R}$ | $R_{eq} = \sum_{j=1}^n R_j$ | $\frac{1}{R_{eq}} = \sum_{j=1}^n \frac{1}{R_j}$ | |
| $q = CV(1 - e^{-t/RC})$ | $i = \frac{dq}{dt} = \frac{V}{R} e^{-t/RC}$ | $q = q_0 e^{-t/RC}$ | $i = \frac{dq}{dt} = -\frac{q_0}{RC} e^{-t/RC}$ | | |
| $F_B = q\vec{v} \times \vec{B}$ | $qvB = \frac{mv^2}{r}$ | $F_B = i\vec{L} \times \vec{B}$ | $dF_B = id\vec{L} \times \vec{B}$ | $\tau = \vec{\mu} \times \vec{B}$ | |
| $\mu = NiA$ | $\epsilon_L = -L \frac{di}{dt}$ | $L = \frac{N\Phi_B}{i}$ | $d\vec{B} = \frac{\mu_0 i}{4\pi} \frac{d\vec{s} \times \vec{r}}{r^3}$ | | |
| $\oint \vec{B} \cdot d\vec{s} = \mu_0 i_{enc}$ | $B = \mu_0 in$ | $B = \frac{\mu_0 i}{2\pi r}$ | $\Phi_B = B \cdot A$ | | |
| $\epsilon = -\frac{d\Phi_B}{dt}$ | $\epsilon = v\ell B$ | $V_s = V_p \frac{N_s}{N_p}$ | $I_s = I_p \frac{N_p}{N_s}$ | | |
| $I = \frac{\mathcal{E}_m}{Z}$ | $V_{rms} = \frac{\mathcal{E}_m}{\sqrt{2}}$ | $X_L = \omega_d L$ | $X_C = \frac{1}{\omega_d C}$ | $Z = \sqrt{R^2 + (X_L - X_C)^2}$ | $\tan \phi = \frac{X_L - X_C}{R}$ |